

INDEX

COPEIA, 1979, Nos. 1-4

NEW NAMES

- Cophixalus concinnus* 118-121
Cynisca rouxae 122-125
Diceratias pileatus 140-141
Etheostoma blennius sequatchiense 198-201
Kinosternon flavescens durangoense 219-220
Libotonius pearsoni 400-405
Microdesmus africanus 203-205
Phrynychthys thele 142-143
Plethodon fourchensis 109
Rana chiricahuensis 383-390
Rhabdosargus thorpei 702-707
Rypticus courtenayi 396-398
Siganus puelloides 390-393
Telmatobius cirrhacelis 715-721
Umbrivaga pyburni 698-701
Urolophus tumbesensis 709-713
Varicus marilynae 126-128

NEW AND ESTABLISHED NAMES

- Abramis brama* 27
Abudefduf troschelii 242-249
Acanthopagrus berda 268
Acarichthys heckeli 679-691
Acipenser
 ruthenus 518
 stellatus 524
Acrochordus javanicus 744-745
Adelotus brevis 297-306
Aelurophryne
 (2 spp.) 297-306
 brevipes 297-306
Aeluroscalabotes 9
Aequidens
 metae 679-691
 paraguayensis 679-691
Afrixalus pygmaeus 278-285
Afroedura transvaalica platiceps 15-16
Agalychnis
 annae 650-655
 callidryas 650-655
Agama agama 585-619
Agirodon piscivorus 460, 549
Alligator mississippiensis 48-59, 549
Alvordius 413-426
 crassus 421
 roanoka 416
Amblyrhynchus cristatus 255, 496
Ambystoma
 (5 spp.) 297-306
 gracile 273
 maculatum 270-274, 348-350
 tigrinum 348-350, 747-749
Amia calva 404
Amphibolurus
 caudicinctus 38
 decrepii 29-40
 fionii 29-40
 ornatus 38
 rufescens 38
 vadnappa 29-40
Amphilophus 679-691
Amphiplaga 400-401, 404
 brachyptera 401, 403
Amphisbaena 125
Amphiuma tridactylum 297-306
Amyzon 404
Anarchias yoshiae 128
Anableps
 anableps 88-89
 dowii 84
 dowei 84
 dowi 82-91
 dowii 84
 microlepis 83-84, 88-89
Aneides arneus 332, 654
Anguilla
 anguilla 360
 japonica 362
 rostrata 360-363
Anodonthyla boulengeri 653-654
Anniella
 geronimensis 242-249
 pulchra 242-249
Anolis 14, 19, 150
 acutus 242-249, 580
 aeneus 242-249
 bimaculatus 242-249
 biporcatus 481
 blanquillanus 242-249
 bonaerensis 242-249
 carolinensis 432, 482, 623
 chlorocyanus 482
 chochorum 481
 coelestinus 482
 cristatellus 243, 482
 cybotes 482
 extremus 242-249
 garmani 482
 gingivinus 242-249
 grahami 481-494
 griseus 242-249
 hendersoni 482
 humilis 472-481
 isolepis 482
 leachi 242-249
 limifrons 472-481, 620-626
 lionotus 488, 490
 lividus 242-249
 luciae 242-249
 lucius 482
 marmoratus 242-249

- Chelys fimbriatus* 657
Chiromantis 633
 petersii 278-285
Chiropterotriton magnipes 741-744
Choeophryne 118
Chondrodactylus 14
Chriolepis 126
 benthonis 126
 minutillus 126
 new species 126, 128
Chrysemys 697
 decorata 692-698
 floridana 659
 picta 165-169, 551, 659
 scripta 168, 549
Chrysophrys
 aries 707
 chrysargyra 707
 globiceps 702, 708
 natalensis 705, 707
Cichla
 ocellaris 43-47
 temensis 679-691
Cichlasoma 679-691
 beani 679-691
 bimaculatum 679-691
 centrarchus 679-691
 citrinellum 543, 679-691
 coryphaenoides 679-691
 cyanoguttatum 679-691
 dovii 543, 679-691
 dowi 543, 679-691
 facetum 686
 festivum 679-691
 kraussi 679-691
 labridens 679-691
 macracanthum 85
 maculicauda 46
 managuense 679-691
 meeki 679-691
 motaguense 85
 nigrofasciatum 85, 679-691
 octofasciatum 679-691
 salvini 679-691
 septemfasciatum 679-691
 severum 679-691
 sp. 679-691
 spilurum 680
 trimaculatum 85, 679-691
Citharichthys spilopterus 535-538
Clarias
 batrachus 156-158
 gariepinus 268
Clemmys guttata 165-169
Clinostomus elongatus 26
Cnemaspis 553
Cnemidophorus
 murinus 573-577
 sexlineatus 747-749
 tigris 242-249, 576, 580
Cnidogobius macrocephalus 358-359
Colisa fasciatus 146-149
Colopus 14
Colostethus
 collaris 297-306
 trinitatus 297-306
Constrictor constrictor 53
Cophixalus
 concinus 118-121
 exiguus 119, 121
 neglectus 119, 121
 ornatus 119, 121
 saxatilis 119, 121
Cophosaurus 36
Copiula 118
Coryphaenoides
 macrocephalus 541-542
 thelestomus 541-542
 zaniophorus 541-542
Cottus pollus 148
Crenadactylus ocellatus 13
Crenicara filamentosa 679-691
Crenicichla
 lepidota 679-691
 lucius 679-691
 notophthalmus 679-691
 strigata 679-691
Crocodylus
 acutus 48-59, 549
 niloticus 55, 162, 266-269
Crotalus
 atrox 460
 lepidus 638-639, 641-642
Crotaphytus collaris 580
Cryptobranchus alleganiensis 92-95, 297-306
Ctenosaura pectinato 431-432
Culaea inconstans 410
Cyclorana dahlui 297-306
Cynisca
 bifrontalis 124-125
 degrysi 123-125
 feae 124-125
 gansi 124-125
 haughi 124-125
 kigomensis 124-125
 kraussi 123-125
 lamottei 124-125
 leonina 124-125
 leucura 124-125
 liberiensis 124-125
 mulleri 124-125
 nigeriensis 124-125
 oligopholis 124-125
 rouxae 122-125
 schaeferi 124-125
Cynoscion regalis 535-538
Cyprinodon 149-152
 alvarezi 149-152
 alvarezi × *Megupsilon aporus* 149-152
 eximius 149
 rubrofluvialis 79
Cyprinus carpio 451
Cyrtodactylus 6, 12

Damalichthys vacca 567-572
Deirochelys insculpta 656, 659

- Dendrobates*
(14 spp.) 297-306
auratus 297-306
lehmanni 297-306
- Desmognathus*
(7 spp.) 297-306
fuscus fuscus 354
monticola 334
ochrophæus 270-274, 332-333, 650, 654
quadramaculatus 273
- Diadophis punctatus* 334, 460
- Diaphus urolampus* 555
- Diceratias* 129, 138-139, 141, 143
bispinosus 129, 132, 134, 136, 139-141, 143
pileatus 129, 141, 143-144
- Dicroglossus occipitalis* 346-348
- Diplodactylinae (subfamily of Gekkonidae) 1-21
- Diplodactylus* 6, 15-16
byrnei 11, 13
ciliaris 11, 13
conspicillatus 11, 14
damaeus 10-11, 13-14
elderi 11, 13
pulcher 11
spinigerus 11
steindachneri 11
stenodactylus 11
strophurus 11
tesselatus 11
vittatus 11, 14
- Diplodus sargus* 707
- Dipsosaurus dorsalis* 55-56, 255, 262-263, 431-432, 435, 576, 580, 585, 592, 594, 599
- Discodeles bufoniformis* 346-348
- Dromicus* 701
- Drymarchon corais* 329-341
- Ebenavia inunguis* 12-13
- Egernia cunninghami* 431-432, 435
- Eigenmannia* 752
- Elaphe* 460
guttata 319-331, 739
guttata emoryi 319-331
guttata guttata 319-331, 739-741
guttata rosacea 319-331
longissima 329
obsoleta 319-331
obsoleta obsoleta 319-331
obsoleta quadrivittata 319-331, 739
obsoleta spiloides 319-331
quatuorlineata 529
vulpina 319-331
vulpina vulpina 319-331
- Elapomorphus*
bilineatus 745-747
bollei 745
iheringi 745
lemniscatus 746
melanopleurus 745
reticulatus 745
spgazzinii 745-746
suspectus 745-746
tricolor 745
trilineatus 745
- Electrophorus* 752
- Eleutherodactylus latrans* 765
- Embiotoca*
jacksoni 571
lateralis 567-572
- Emydoidea* 656
- Enhydryna schistosa* 307-318
- Enhydryn bennetti* 671
- Ensantina eschscholtzi(i)* 297-306, 339, 502-508, 743
- Eohiodon*
rosei 404
woodruffi 400, 404
- Eosalmo* 404
driftwoodensis 404
- Erismatopterus* 400-401, 404
levatus 401-403
- Erythrolamprus aesculapii* 169-172
- Etheostoma*
blennioides 196-197, 199, 202
blennius 191-203
blennius: blennius × *sequatchiense* 191-203
blennius blennius 191-203
blennius sequatchiense 191-203
euzonum 199
histrio 197, 199, 202
inscriptum 197, 199
kanawhae 199
luteovinctum 196
nigrum 426-427, 429
nigrum eulepis 428
nigrum: nigrum × *susanae* 426-430
nigrum nigrum 426-427, 429
nigrum susanae 426-430
osburni 199
peltatum crassus 421
roanoka 416
rupestre 197, 199
sellare 197, 199
spp. 195
striatulum 196
swannanoa 196-197, 199, 202
tetrazonum 197, 199
thalassinum 196-197, 199
variatum 196-197, 199
zonale 196-197, 199, 202
- Eucyclogobius newberryi* 530
- Eumeces*
anthracinus 748
fasciatus 748
laticeps 748
obsoleta 580, 747-749
septentrionalis 747-749
- Euproctus*
asper 297-306
montanus 297-306
- Eurycea*
bislineata 270-274
longicauda 297-306
- Fordonia leucobalia* 671
- Fundulus*
diaphanus 544-546
heteroclitus 291-297, 358-359
majalis 291

notat
spp.

Galeich
Gambu
affin
affin
affin
affin
Garmar
Gastero
acule
acule
acule
acule
Gastroth
Gaviadi
Gehyra
mutil
Gekko
Gekko
Geochel
dentia
elong
Geophag
brasil
jurup
surin
Gerrhon
Gerrhos
Gillichth
Girella
Gnathar
Gnathop
Gobiom
Gopheru
Gyrinop

Hadrop
Heleiop
(3 sp
cyrei
Hemida
flaviu
frenae
garno
Hemierg
Hemiph
Hemiph
Hemisus
Herichth
Heros 6
Herotila
Heterod
nasicu
platyr
Hildatia
brasil
Hildebra
Hipopom
Holacan
bermu
ciliaris
tricolor

- notatus* 291
spp. 152
- Galeichthys felis* 359
- Gambusia*
affinis 148, 452, 509-513
affinis: affinis × *holbrooki* 509-513
affinis affinis 60-64, 509-513
affinis holbrooki 509-513
- Garmanella pulchra* 152
- Gasterosteus* 405, 411
aculeatus 525-535, 635-638
aculeatus aculeatus 529
aculeatus microcephalus 529-533
aculeatus williamsoni 529, 532
- Gastrotheca peruana* 297-306
- Gavialis gangeticus* 162
- Gehyra* 17
mutilata 552-554
- Gekko* 17-18
- Gekkoninae (subfamily of Gekkonidae) 1-21
- Geochelone*
denticulata 697
elongata 659
- Geophagus*
brasiliensis 679-691
jurupari 679-691
surinamensis 679-691
- Gerrhonotus multicarinatus* 580
- Gerrhosaurus* cf. *major* 172-174
- Gillichthys mirabilis* 63
- Girella nigricans* 62
- Gnathanodon speciosus* 535-538
- Gnathopogon coerulescens* 22-28
- Gobiomorus dormitor* 542-544
- Gopherus polyphemus* 659
- Gyrinophilus porphyriticus* 270-274, 334
- Hadropterus roanoka* 416
- Heleioporus* 303
(3 spp.) 297-306
cyrei 297-306
- Hemidactylus* 6, 17
flaviviridis 553
frenatus 552-554
garnoti 553
- Hemiergis peronii* 66
- Hemiphractus panamensis* 297-306
- Hemiphyllodactylus* 17
- Hemistis marmoratum* 278-285
- Herichthys* 679-691
- Heros* 679-691
- Herotilapia multispinosa* 679-691
- Heterodon*
nasicus 748
platyrhinus 739, 747-749
- Hildatia* 751-752
brasiliensis 751-752
- Hildebrandtia ornata* 279
- Hipopomus* 752
- Holacanthus*
bermudensis 161
ciliaris 161
tricolor 160-162
- Holbrookia* 36
maculata 747-749
- Holocentrus rufus* 161
- Hoplosternum* 157
- Hydromantes* 442, 446
genei 297-306, 445
- Hydrophis* 671-672
belcheri 494-497
cyanocinctus 494-497
fasciatus 316
ornatus 551
- Hyla*
(13 spp.) 297-306
bifurca 626-635
cruentomma 626-635
eberracata 650-655
faber 297-306
garbei 626-635
gratiosa 297-306
pardalis 297-306
parviceps 626-635
punctata 626-635
regilla 297-306
rhodopepla 626-635
rubra 626-635
sarayacuensis 626-635
triangulum 626-635
versicolor 286-290
- Hylidae*
(4 Asian spp.) 297-306
(41 Australian spp.) 297-306
(1 North American sp.) 297-306
(135 South American spp.) 297-306
- Hymenochirus boettgeri* 297-306
- Hynobius*
(2 spp.) 297-306
nasutus 297-306
- Hyperolius*
marmoratus 297-306
nasutus 297-306
- Hypopomus* 752
- Icichthys* 733, 735
lockingtoni 735
- Ictalurus*
melas 452
nebulosus 359
- Idiolychnus urolampus* 555
- Iguana iguana* 497
- Jordanella floridae* 152
- Kassina senegalensis* 278-285
- Kathetostoma albigutta* 128
- Kinosternon*
arizonense 175-177, 218, 221
bauri(i) 548-552, 659, 692-698
baurii baurii 548-552
baurii palmarum 551
cruentatum 219
flavescens 175, 212-225
flavescens arizonense 175-177, 212-225
flavescens durangoense 212-225
flavescens flavescens 175-176, 212-225

- flavescens: flavescens* × *durangoense* 217, 224
flavescens spooneri 212-225
flavescens stejneri 175-176, 219-221
herrerai 218-219
hirtipes 175, 218-219
integrum 175, 218-219
scorpioides 218-219
scorpioides abaxillare 692-698
scorpioides cruentatum 692-698
scorpioides scorpioides 693
sonoriense 175, 218-219
spurrelli 549-551
subrubrum 218-219
subrubrum steindachneri 692-698
Klauberina riversiana 431-437, 577-584
Kyarranus kundagungan 297-306
- Lacerta**
dugesii 250-257
dugesii mauli 250
trilineata 580
vivipara 22, 432
Laevoceratias liparis 138
Lampropeltis
calligaster 747-749
getulus 748
getulus holbrookii 329, 642
triangulum syspila 329
Lapemis hardwickii 316, 496
Laticauda 307, 551, 661, 672
colubrina 494-497, 672
laticaudata 494-497
Leimadophis 698, 701
poecilogyrus 169-172
pygmaeus 698, 700-701
reginae 169-172
Leiocephalus 36
carinatus coryi 36
Leiolopisma
laterale 580
metallicum 66
sp. 66
trilineatum 66
Lepidodactylus 17
lugubris 553
Lepidophyma
gaigeae 577-584
smithii 436, 577-584
Lepisosteus tropicus 84
Lepomis
cyanelus 155-156
gibbosus 635-638
spp. 452
Leptocottus armatus 530
Leptodactylidae
 (37 Australian spp.) 297-306
 (6 North American spp.) 297-306
 (21 South American spp.) 297-306
Leptodactylus
 (8 spp.) 297-306
pentadactylus 303
rugosus 297-306
Leptodeira annulata polysticta 744
- Leptopelis**
bocagei 279
concolor 279
Lerista bougainvillea 66
Leuresthes
sardina 79
tenuis 291, 295
Libotoniis
blakeburnensis 400-404
pearsoni 400-405
Limnodynastes
dumerili 297-306
fletcheri 297-306
peronii 297-306
Lineatriton lineola 742
Liolaemus
fuscus 546-548
lemniscatus 546-548
monticola 546-548
nigromaculatus 546-548
nigroviridis 546-548
nitidus 546-548
platei 546-548
schoederi 546-548
Liophis 698, 701
Litoria aurea 355-356
Lucasium damaeum 14
Lygophis 701
Lymnodynastes sp. 66
Lythrypnus nesiotes 128
- Malaclemys terrapin* 548-549, 551-552, 659, 668
Malacocheilus tornieri 659
Megapsilon aporus 149-152
Melaniris guija 85
Melanostigma
atlanticum 146, 363-366
pammelas 366-369
Menetia greyi 66
Mesonauta 679-691
Microdesmus
aethiopicus 203, 205
affinis 205
africanus 203-205
carri 205
longipinnis 203-205
suttkusi 205
Microhylidae (26 spp.) 297-306
Micropterus salmoides 149
Micruroides euryxanthus 172
Micrurus
corallinus 169-172
fulvius 172, 453-463
fulvius fulvius 453, 458-459
fulvius tener 453-463
Misgurnus fossilis 27
Mogrundera obscura 148
Monacanthus hispidus 535-538
Morethia lineocellatus 66
Morone
americana 513-518
americana × *Morone saxatilis* 513-518
chrysops 513-518

- chrysops* × *Morone saxatilis* 513–518
saxatilis 513–518
Moxostoma
 macrolepidotum 152–154
 macrolepidotum breviceps 154
 macrolepidotum: *macrolepidotum* × *breviceps* 154
Mugil cephalus 266–269
Muraena retifera 128
Muraenesox bagio 268
Mystus tengara 148

Nannacara anomala 679–691
Natrix
 rhombifera 465, 496
 sipedon 452
 taxispilota 53
Necturus (2 spp.) 297–306
Neotrophus nematopus 543, 679–691
Neoceratodus 116
Nephrurus 6, 9–10, 14
 asper 9
 laevissimus 10
 levis 9–10
Nerodia
 fasciata 671
 fasciata clarki 548, 551
 fasciata compressicauda 548–549
 fasciata taeniata 548
 rhombifera 465
 sipedon 551, 671
 sipedon williamengelsi 548
Nocomis micropogon 429
Notophthalmus
 meridionalis 297–306
 viridescens 274, 297–306, 351
Notropis
 girardi 77
 lutrensis 70–81
 volucellus 437–441
Nyctimystes
 humeralis 297–306
 rueppeli 297–306

Oedipina (7 spp.) 297–306
Oedomognathus exodon 751–752
Oedura 6, 15–17
 castelnaui 15
 lesueurii 15
Oneirodes 129, 136, 143
Onychodactylus japonicus 297–306
Ostariophysan fishes 111–118
Oxyrhopus petola digitalis 169–172
Oxyzygonectes dowi 87

Pachydactylus 14
Pachymedusa dactinolor 297–306
Paraliparis
 calidus 145–146
 copei 145–146
 garmani 145–146
Paramesotriton chinensis 297–306
Parapetenia 679–691
Paroedura gracilis 12–13

Paroneirodes 129, 139, 141
 glomerosus 129, 139, 141–142
Pelamis platurus 307, 672
Pelobatidae
 (11 Asian spp.) 297–306
 (3 North American spp.) 297–306
Pelophryne misera 297–306
Peprilus
 burti 538–541
 burti × *Peprilus triacanthus* (?) 538–541
 triacanthus 538–541
Perca flavescens 635–637
Percina
 crassa 413–426
 crassa roanoka 416
 notogramma 413–416, 420–425
 notogramma montuosa 420
 notogramma notogramma 420
 pellata 413–426
 roanoka 413–426
Percopsis 401, 403–404
 omiscomaycus 403
 transmontana 403
Phoxinus cumberlandensis 426
Phrynichthys 129, 141, 143
 sp. 141–143
 thele 129, 139–143
 wedli 129–144
Phrynobatrachus
 acridoides 278–287
 natalensis 278–287
Phrynohyas (3 spp.) 297–306
Phrynomerus bifasciatus 278–287
Phrynosoma 262
 cornutum 747–749
 douglasi 747–749
 m'calli 580
 spp. 263
Phyllobates
 (5 spp.) 297–306
 zaparo 297–306
Phyllodactylus 6, 16
 europaeus 12–13
 gerrhopygus 12
 julieni 12
 marmoratus 12
 microphyllus 12
 palmatus 12
 porphyreus 12, 553
 riebeckii 13
 siamensis 12
Phyllurus cornutus 9
Physalaemus
 lemur 650–655
 pustulosus 650–655
Pimphales promelas 451, 549
Pipidae (4 spp.) 297–306
Pituophis melanoleucus melanoleucus 329
Placogaster 125
Plagiotremus spp. 160–161
Platyhyla grandis 653–654
Platythyra flavescens 218
Platyurus platyurus 552–554

Plethodon

- (9 spp.) 297-306
caddoensis 95-110, 502-508
cinereus 270-274, 332, 339, 502-508, 653
dorsalis 502-508
dunni 502-508
elongatus 502-508
fourchensis 95-110, 502-508
fourchensis × *Plethodon ouachitae* 97, 100, 103-104, 107
glutinosus 95-110, 270-274, 502-508
hoffmani 502-508
hubrichti 502-508
idahoensis 502-508
jordani 107-108, 270-274, 502-508
larseli 502-508
neomexicanus 502-508
nettingi 502-508
ouachitae 95-110, 502-508
punctatus 502-508
richmondi 271, 502-508
serratus 106, 502-508
shenandoah 502-508
stormi 502-508
vandykei 502-508
vehiculum 502-508, 743
websteri 502-508
wehrlei 502-508
welleri 271, 502-508
yonahlossee 95-110, 502-508
Plethodontohyla notosticta 653
Pleurodeles waltl 297-306
Plotosus
anguillaris 359
lineatus 357-360
Poecilia
reticulata 225-242, 440
sphenops 148
spp. 85
Poeciliopsis 85
occidentalis 60
Polyodon spathula 518, 521, 524
Pomadasys commersoni 268
Pranesus insularum 158-159
Prionosparus 702
tricuspidens 702
Promicropterus decoratus 396, 398
Protopterus amphibius 360
Pseudemys (= *Chrysemys*) *scripta* 549
Pseudoeurycea juarezi 742
Pseudophryne 654
(3 spp.) 297-306
corroboree 303
Pseudopleuronectes americanus 358-359
Pseudoscaphirhynchus kaufmanni (= *kaufmanni* ?) 524
Pseudotomodon trigonatus 745
Pseudotriton
montanus 271, 274
ruber 271, 274, 334
Pterophyllum
eimekei 679-691
scalare 679-691
Ptychadena superciliaris 278-287

Pungitius

- hellenicus* 405, 411
platygaster 405, 408-409, 411
platygaster aralensis 405, 411
platygaster platygaster 405, 408-409, 411
pungitius 405-412
pungitius larvis 405-409, 411
pungitius pungitius 405-409, 411
pungitius sinensis 405, 411
pungitius tymensis 405, 411

Python

- molurus* 56, 466-471
molurus bivittatus 466-471
molurus molurus 466-471

Pyxicephalus

- adspersus*(a) 297-306, 346-348
delaland(e)i 278-287, 346-348

Rana 40

- (10 Asian spp.) 297-306
(3 North American spp.) 297-306
albolabris 346-348
angolensis 346-348
areolata circulosa 346-348
arfaki 346-348
arvalis 346-348, 554
aurora 346-348
berlandieri 385, 389, 643-650
berlandieri forreri 387
blairi 342, 385-386, 389
blythi 346-348
boylei 346-348
cancrivora 346-348
cascadeae 346-348
catesbeiana 302-303, 346-348, 447-453, 554
chiricahuensis 385-390
chiricahuensis × *Rana magnaocularis* 389
daemeli 346-348
dunni 346-348
erythraea 346-348
esculenta 346-348, 554
fisheri 388-389
fuscigula 346-348
galamensis 346-348
grisea 346-348
gryllio 346-348
heckscheri 346-348
hexadactyla 346-348
johni 346-348
kuhlii 297-306
lessonae 554
maculata 346-348
magnaocularis 385, 389
megapoda 346-348
montezumae 346-348
muscosa 346-348
narina 346-348
onca 388-389
palmipes 346-348, 554-555
palustris 346-348
papua 346-348
pueblae 346-348
pustulosa 346-348

ridi
sept
sier
sph
sylo
tara
temp
tigr
utris
vibi
virg
zenk
Ranid
(62
(21
Regin
sept
Rhadr
glob
holu
sarb
thor
tricu
Rhaco
auri
Rhaco
Rhadin
Rhadin
guat
Rhino
Rhino
sp. 6
Rhynch
Rhynch
macu
nigri
xanti
Roeboi
Roughl
Rutilus
Rypticu
bicol
bistr
court
macu
nigri
sapon
sapon
sapon
Sacalia
Salama
(2 sp
salam
Salama
Salmo
clark
gaird
trutta
Sanzin
Sargus
auriv
holub

- ridibunda* 554
septentrionalis 346-348
sierramadrensis 346-348
sphenocephala 643
sylvatica 346-348
tarahumarae 346-348
temporaria 346-348, 554
tigrina 346-348
utricularia 346-348, 385, 643
vibicaria 346-348
virgatipes 346-348
zenkeri 346-348
- Ranidae**
 (62 Asian spp.) 297-306
 (21 North American spp.) 297-306
- Regina* 465
septemvittata 551
- Rhabdosargus* 702
globleps 702-709
holubi 702-709
sarba 702-709
thorpei 702-709
tricuspidens 702-709
- Rhacodactylus* 6, 16-18
auriculatus 16, 18
- Rhacophoridae* (48 African spp.) 297-306
- Rhadinaea* 698
- Rhamdia* 86
guatemalensis 85
- Rhinobatos annulatus* 268
- Rhinoclemmys* 697
 sp. 659
- Rhynchoedura ornata* 13
- Rhypticus*
maculatus 398
nigripinnis 398
xanti 395
- Roeboides* 85
- Roughleyia taruhine* 707
- Rutilus rutilus* 27
- Rypticus*
bicolor 393-400
bistripinis 399
courtenayi 393-400
maculatus 396, 398
nigripinnis 393-400
saponaceus 393, 396
saponaceus bicolor 393, 396
saponaceus saponaceus 393
- Sacalia* 697
- Salamandra*
 (2 spp.) 297-306
salamandra 442-447
- Salamandrina* (2 spp.) 297-306
- Salmo*
clarki 673-678
gairdneri 635
trutta trutta 534
- Sanzinia madagascarensis* 470
- Sargus*
auriventris 702, 707
holubi 702, 707-708
- natalensis* 708
nigrofasciatus 708
- Sauromalus obesus* 255-256, 431-432, 435-436
- Scaphiopus (Spea) bombifrons* 747-749
- Scaphirhynchus platyrhynchus* 518-525
- Scarus* 211
blochii 206-212
capistratoides 206-212
croicensis 206-212
fasciatus 206-207
gibbus 207, 209
inserti 206-212
japanensis 206-212
jonesi 207
perspicillatus 207, 210
quoyi 206-212
rhoduropterus 207
sordidus 207-210
taeniopterus 211
trocheli 207
viridis 206-207, 209, 211
- Sceloporus*
cyanogenys 258-266
graciosus 432, 460, 580, 624
jarrovi 624
occidentalis 263, 432, 576
olivaceus 580
torquatus 37
undulatus 168, 747-749
undulatus garmani 748
- Schedophilus*
griseolineatus 735
huttoni 733, 735-736
labyrinthicus 735
medusaphagus 733, 735-736
- Scincella lateralis* 54
- Sclerophages* 116
- Scutigera pingii* 297-306
- Scymnus* 524
- Semotilus atromaculatus* 26
- Seriotelella* 735
- Siebenrockiella* 697
- Siganus*
argenteus 390
corallinus 390
javus 390
lineatus 390
luridus 390
magnificus 390
puelloides 390-393
puellus 390, 392
rivulatus 390
stellatus 390
sutor 390
trispilos 390
vulpinus 390
- Sistrurus*
miliarius 638-642
miliarius barbouri 638-642
miliarius streckeri 638-642
- Sparisoma viride* 206-207
- Sparus*
bufonites 707

- psittacus* 707
sarba 702, 707
Sphenocephalus 404
 spp. 255
 quoyi 261, 263
Sphyraena barracuda 159
Spinachia spinachia 410
Stauroids natalor 346-348
Staurotypus
 savlinii 692-698
 triporatus 693
Stenodactylus 14
 khobarensis 9-10
 slevini 9
Stereochilus marginatus 297-306
Sternopygus 751-752
 brasiliensis 752
Sternotherus
 minor minor 692-698
 odoratus 165-169, 455, 551, 655-660
Stolephorus purpureus 158-159
Storeria dekayi 463-466
Strongylura marina 535-538
Symphysodon aequifasciata 679-691

Tarentola 553
Taricha torosa 297-306
Telmatobius
 brevipes 732
 cinereus 714, 724-725
 cirrhacelis 714-733
 culeus 714, 731
 ignavus 731
 latirostris 732
 niger 714-733
 vellardi 714-733
Terrapene
 carolina 655-660
 ornata 53, 168
Thalassophis anomalus 316
Thamnophis 307
 sirtalis 329, 464
 sirtalis sirtalis 329
Thorichthys 679-691
Tiliqua 262
Trachurus lathami 535-538
Trichogaster labiosus 148
Trichophanes 404
Trinectes 86
Trionyx
 ferox 659
 spiniferus 655-660
Triturus
 (9 spp.) 297-306
 alpestris 40-43
 cristatus 40, 350-353
 vulgaris 40, 350-351
Tropidurus spp. 38

Trygonoptera 709
Tubbia tasmanica 733-738
Tylotriton (2 spp.) 297-306
Typhlops braminus 744

Uaru amphicanthoides 679-691
Umbrivaga
 mertensi 698-701
 pyburni 698-701
 pygmaeus 698-701
Urolophus
 armatus 713
 bucculentus 712
 circularis 712
 concentricus 709-713
 halleri 709-713
 jamaicensis 709-713
 javanicus 712
 kaianus 712
 maculatus 709-713
 muscosus 712
 paucimaculatus 713
 testaceus 712
 tumbesensis 709-713
 viridis 713
Uromastix aegyptius 432
Urosaurus spp. 36
Urotheca 698
Urotrygon 709
Uta
 mearnsi 580
 stansburiana 256, 580, 612, 624

Varanus
 bengalensis 67-69
 brevicauda 69
 caudolineatus 69
 eremius 69
 exanthematicus 65
 gouldii flaviviridis 64-65, 67-69
 gouldii gouldii 64, 68
 gouldii rosenbergi 64-70
 mertensi 69
 "monitor" 67
 niloticus 68-69
 spenceri 68-69
Varicus
 bucca 126-128
 marilynae 126-128
 sp. 126
Vipera ammodytes 642

Xantusia
 henshawi 577-584
 vigilis 242-249, 455, 577-584
Xenodermichthys copei 363-366
Xenodon sp. (probably *merremii*) 169-172
Xenopus 750

Ypsiscarus oedema 207, 209

SUBJECT INDEX

- ABUNDANCE, 13 anuran species (distribution and abundance in Kenyan pond) 278-285.
- ACOUSTICS, *Rana berlandieri* (acoustic behavior of males: experimental analysis through field playback trials) 643-650.
- ACTIVITY, *Cryptobranchus alleganiensis* (diel activity rhythms) 92-95; *Liolaemus* (8 spp.) (activity temperatures) 546-548.
- ADAPTATION, *Cichla ocellaris* (adaptive significance of color patterns) 43-47; *Desmognathus ochrophaeus* (adaptiveness of parental care of egg clutches) 332-341.
- ANATOMY, two subfamilies of gekkonine and dipodactyline geckos (feet) 1-21; *Rana palmipes* (blood vascular system) 554-555.
- AUDITION, *Anolis grahmi* (behavioral demonstration) 490-494.
- BEHAVIOR, *Amphibolurus decresii* species complex (hind-leg pushup display) 29-40; *Triturus alpestris* (significance of color signals in partner recognition) 40-43; *Alligator mississippiensis* and *Crocodylus acutus* (with temperature changes) 48-59; snakes (tail displays in response to predation by mustelid mammals (genus *Galictis*)) 169-172; *Desmognathus ochrophaeus*, *Eurycea bislineata*, *Plethodon cinereus*, *jordani*, *glutinosus*), *Ambystoma maculatum*, *Gyrinophilus porphyriticus* (effectiveness of behavior and secretions against shrew predation) 270-274; *Elaphe* (*obsoleta*, *guttata*, *vulpina*) (reproductive behavior) 319-331; *Desmognathus fuscus* (larval feeding behavior) 354; *Salamandra salamandra* (relationship between stimulus orientation and stimulus movement in prey catching behavior) 442-447; *Python molurus* (social behavior) 466-471; *Anolis* (*humilis*, *limifrons*) (inter- and intraspecific aggression) 472-481; *Anolis grahmi* (learning paradigm and behavioral demonstration of audition) 490-494; *Agama agama* (microclimate relationships) 585-593; *Agama agama* (implications of mechanistic ecology) 594-614; *Sistrurus miliarius* (combat ritual) 638-642; *Rana berlandieri* (acoustic behavior of males: experimental analysis through field playback trials) 643-650.
- BIOCHEMISTRY, *Anguilla rostrata* (phosphate compounds in red cells of liver and skeletal muscle) 360-363; *Carassius auratus* 749-751.
- BLOOD, *Anguilla rostrata* (phosphate compounds in red blood cells) 360-363; sea snakes (4 spp.) (blood lactate in free-diving snakes) 494-497; *Rana palmipes* (anatomy of blood-vascular system) 554-555.
- BREEDING, *Hyla versicolor* (mate selection related more to perch sites rather than size of male; relation to breeding success) 286-289; *Elaphe* (*obsoleta*, *guttata*, *vulpina*) (reproductive behavior) 319-331; *Gasterosteus aculeatus* (nest habitat preference of low-plate number morphs) 525-528; *Gasterosteus aculeatus* (low-plate number morph breeding in salt water) 529-533; *Gobiomorus dormitor* (breeding biology) 542-544.
- CALLING, *Rana berlandieri* (acoustic behavior) 643-650.
- CHROMOSOMES, *Gambusia affinis* (sex chromosome evolution) 509-513; *Fundulus diaphanus* (location of nucleolus organizer regions of sex chromosomes) 544-546; neotropical Cichlidae (41 spp.) (cytotaxonomy) 679-691; genera *Kinosternon*, *Sternotherus*, *Staurotypus* (6 taxa) (banded karyotypes) 692-698.
- COLOR, *Triturus alpestris* (significance of color signals in partner recognition) 40-43; *Cichla ocellaris* (adaptive significance of color patterns) 43-47; *Poecilia reticulata* (genetic studies of melanistic color patterns) 225-239; *Poecilia reticulata* (melanophore-based color mutants; cytological aspects and differential response to melatonin) 232-242; *Lacerta dugesii* (dorsal pigmentation determined to be related to selective predation pressures rather than thermal regulation) 250-257.
- COMBAT, *Sistrurus miliarius* (combat ritual) 638-642.
- COURTSHIP, *Triturus alpestris* (significance of color) 40-43.
- CYCLES, *Fundulus heteroclitus* (lunar spawning cycle) 291-297.
- CYTOLOGY, *Poecilia reticulata* (cytological aspects and differential response to melatonin of melanophore-based color mutants) 232-242.
- DEMOGRAPHY, *Gasterosteus aculeatus* (population decline and change in frequency of lateral plates) 635-638.
- DENTITION, *Gnathopogon coeruleus* (dentition; development and replacement pattern of pharyngeal teeth in Japanese cyprinid fish) 22-28.
- DEVELOPMENT, *Gnathopogon coeruleus* (of pharyngeal teeth) 22-28; *Chiropterochilus magnipes*, *Bolitoglossus magnipes* (egg development time) 741-744.
- DIET, *Varanus gouldii rosenbergi* 64-69; genus *Paraliparis* (*copei*, *calidus*, *garmani*) 145-146; *Etheostoma blennioides* 202.
- DIGESTION, *Klauberina riversiana* (digestive efficiency) 431-437; *Scaphirhynchus platyrhynchus* (histology of digestive organs) 518-525.
- DIMORPHISM, numerous amphibian spp. (sexual selection and sexual dimorphism) 297-306; *Gasterosteus aculeatus* (sexual dimorphism in integumentary tissues) 533-535.
- DISEASE, *Centrolenella prosoblepon*, *Agalychnis callidryas*, *annae* (fungi lethal to frog eggs) 650-655.
- DISTRIBUTION, family Diceratiidae (maps) 141; *Paraliparis* (*copei*, *calidus*, *garmani*) (geographical and depth) 145-146; *Etheostoma blennioides* (maps) 202; *Kinosternon flavescens* (4 subsp.; United

States and Mexico) 213; 13 anuran spp. (abundance and distribution in a Kenyan pond) 278-285; *Litoria aurea* (current distribution of introduced populations in New Hebrides) 355-356; *Elapomorphus bilineatus* (southernmost population; Argentina) 745-747.

ECOLOGY, *Notropis lutrensis* (influence of physico-chemical factors on habitat selection) 70-81; *Anableps dowi* (general; comparison with other 2 spp. in genus) 82-91; *Cryptobranchius alleganiensis* (diel activity rhythms) 92-95; *Varicus marilynae* (notes) 126-128; *Paraliparis (copei, calidus, garmani)* (notes) 145-146; *Etheostoma blennioides* (general; spawning; diet) 191-203; *Anolis (humilis, limifrons)* (niche overlap and inter- and intra-specific aggression) 472-481.

EGGS, *Desmognathus ochrophaeus* (egg clutches less susceptible to predation when attended by females than when unattended) 332-341; *Ambylostoma (tigrinum, maculatum)* (ontogenetic variation) 348-350; *Gehyra mutilata*, *Hemidactylus frenatus*, *Platyurus platyurus* 552-554; *Hyla* (9 Ecuadorian spp.) 625-635; *Centrolenella prosoblepon*, *Agalychnis (callidryas, annae)* (fungi lethal to eggs) 650-655; *Chiropoterotriton magnipes*, *Bolitoglossa compacta* (egg development time and clutch size) 741-744.

ELECTROPHORESIS, *Plethodon (ouachitae, caddoensis, fourchensis)* (genetic relationships of Ouachita Mountain spp.) 95-110; *Moxostoma macrolepidotum* (creatine kinase variability) 152-153; 31 spp. of herps and fishes (effects of sample size on genetic distance and heterozygosity estimates in electrophoretic studies) 242-249.

ENERGY, *Hyla* (9 Ecuadorian spp.) (clutch energy partitioning) 625-635.

EVOLUTION, genera *Plethodon* and *Ensatina* (albumin evolution and phylogenetic implications) 502-508; *Gambusia affinis* (sex chromosome evolution) 509-513.

FEEDING, *Alligator mississippiensis* and *Crocodylus acutus* (feeding responses to temperature changes) 48-59; *Varanus gouldii rosenbergi* (diet) 64-72; piscivorous predators and disabled prey 158-160; *Holacanthus tricolor* (possible mucophagy by juveniles) 160-162; *Bufo boreas* (olfactory cues in feeding response) 163-165; *Bufo boreas* (feeding responses elicited by lipid components of prey odors) 275-279; *Desmognathus fuscus fuscus* (larval feeding behavior) 354; *Melanostigma pammelas* (feeding in aquaria) 366-369; *Notropis volucellus* (foraging tactics) 437-441.

FEET, gekkonine and diplodactyline geckos (morphology) 1-21.

FOOD, *Anableps dowi* 82-91; genus *Paraliparis (copei, calidus, garmani)* 145-146; *Holacanthus tricolor* (possible mucophagy by juveniles) 160-162; *Caiman latirostris* (ampullarid gastropods as a possible staple food) 162-163; *Etheostoma blennioides*

202; *Damalichthys vacca*, *Embiotoca lateralis* (geographic patterns of prey utilization) 567-572; *Elaphe guttata guttata* (food consumption) 739-741.

FORAGING, *Notropis volucellus* (tactics) 437-441.

FOSSILS, gerrhosaur (Miocene of Kenya) 172-174; *Kinosternon arizonense* (determined to be senior synonym of *K. flavescens stejnegeri*) 175-177; *Libonotus pearsoni* (new species of Percopidae from Eocene of Washington state) 400-405.

FUNCTION, trionychid turtles (packaging problems of head retraction) 655-660.

FUNGI, *Centrolenella prosoblepon*, *Agalychnis (callidryas, annae)* (two species lethal to frog eggs) 650-655.

GENETICS, *Plethodon (ouachitae, caddoensis, fourchensis)* (relationships) 95-110; *Poecilia reticulata* (genetic studies of melanin color patterns and atypical sex determination) 225-231; *Poecilia reticulata* (cytological aspects and differential response to melatonin of melanophore-based color mutants) 232-242; 31 species of herps and fishes (effects of sample size on genetic distance and heterozygosity estimates in electrophoretic studies) 242-249; *Bufo americanus* (heterozygote deficiencies) 498-502; *Salmo clarki* (genetic similarity of two forms) 673-678; neotropical Cichlidae (41 spp.) (cytotaxonomy) 679-691.

GEOGRAPHICAL LOCALITIES

Alabama, *Etheostoma blennioides* 191-203.

Argentina, *Elapomorphus bilineatus* 745-747.

Arizona, *Kinosternon flavescens arizonense* 175-177; *Rana chirahuensis* 383-390.

Arkansas, *Plethodon (ouachitae, caddoensis, fourchensis, glutinosus)* 95-110.

Atlantic Ocean (western north Atlantic), genus *Paraliparis (coheni, calidus, garmani)* 145-146.

Australia, *Amphibolurus (decreii, fionni, vadnappa)* 29-40; *Varanus gouldii rosenbergi* 64-70; genus *Cophixalus* (including new species, *concinus*) 118-121.

Brazil, seven species of snakes 169-172.

California, *Klauberina riversiana* 431-437.

Chile, *Liolaemus* (8 spp.) 546-548.

Costa Rica, *Anolis (humilis, limifrons)* 472-481.

Delaware, *Fundulus heteroclitus* 291-297.

Ecuador, *Hyla* (9 spp.) 625-635; genus *Telmatobius* 714-733.

Europe (various countries), *Pungitius pungitius* 405-412.

Florida, *Varicus marilynae* 126-128; *Clarias batrachus* 156-158.

Ghana, *Agama agama* 585-619.

Great Britain, *Triturus alpestris* 40-43; *Triturus cristatus* 350-353.

Guatemala, *Anableps dowi* 82-91.

Honduras, *Anableps dowi* 82-91.

Indian Ocean, *Siganus puellus* 390-393.

Ivory Coast, *Cynisca rouxae* (new species) 122-125.

Japan, *Gnathopogon coeruleus* 22-28.

Kansas, Nash local herpetofauna (Pleistocene: Af-
tonian) 747-749.

- (geog-
7-572;
) 739-
41.
2-174;
senior
5-177;
psidae
05.
blems
illdryas,
50-655.
ourchen-
reticulata
ns and
ecilia re-
ntial re-
ed color
and fishes
nce and
phoretic
ozygote
tic simi-
tropical
691.
7.
75-177;
fourchen-
, genus
146.
vadnappe)
0; genus
ncinnus)
481.
elmatobius
pungitius
batrachus
turus cris-
122-125.
cene: Af-
- Kentucky, *Etheostoma nigrum susanae* 426-430; *Rana catesbeiana* 447-458.
Kenya, fossil gerrhosaur 172-174; 13 anuran species 278-285.
Louisiana, *Storeria dekayi* 463-466.
Madeira, *Lacerta dugesii* 250-257.
Malaysia, *Enhydryna schistosa* 307-318.
Maryland, *Hyla versicolor* 286-290.
Mexico, *Anableps dowi* 82-91; *Astyanax mexicanus* 369-371; *Megupsilon aporus* and *Cyprinodon alvarezi* 148-152; *Kinosternon flavescens stejnegeri* 175-177.
New Guinea, genus *Cophixalus* 118-121.
New Hebrides, *Litoria aurea* 355-356.
North Carolina, *Desmognathus ochrophaeus* 332-341; *Percina (crassa, roanoka)* 415-426.
Ohio, *Desmognathus fuscus fuscus* 354.
Oklahoma, *Notropis lutrensis* 70-81; *Plethodon (ouachitae, caddoensis, glutinosus)* 95-110.
Pacific Ocean (eastern), *Rypticus (bicolor, nigripinnis, courtneyi)* 393-400.
Panama, *Anolis limifrons* 620-626.
Peru, *Urolophus tumesens* 709-713.
South Africa, *Mugil cephalus* and *Crocodylus niloticus* 266-269.
South Carolina, *Percina crassa* 415-426.
Tennessee, *Etheostoma blennioides* 191-203; *Etheostoma nigrum susanae* 426-430.
Texas, *Micrurus fulvius tenere* 453-463.
Virginia, *Percina roanoka* 415-426.
Washington, *Libinia (pearsoni)* (Eocene fossil) 400-405; *Gasterosteus aculeatus* 635-638.
West Virginia, *Percina roanoka* 415-426.
Wyoming, *Salmo clarki* 673-678.
Zaire, *Microdesmus africanus* 203-205.
GILLS, *Plotosus lineatus* (histophysiology in relation to osmoregulation) 357-360.
GLANDS, *Cerberus rhynchops* (possible new salt gland) 661-672.
GROWTH, *Enhydryna schistosa* 307-318; *Micrurus fulvius tenere* (in Texas) 453-463; *Elaphe guttata guttata* 739-741.
HABITAT, *Notropis lutrensis* (influence of physico-chemical factors in habitat selection) 70-81; *Gasterosteus aculeatus* (low-plate morph breeding in salt water) 529-533.
HABITS, *Anableps dowi* 82-91.
HISTOLOGY, *Varanus gouldi rosenbergi* (testes) 64-70; *Plotosus lineatus* (gills and dendritic organs) 357-360; *Scaphirhynchus platyrhynchus* (feeding and digestive organs) 518-525.
HYBRIDIZATION, *Plethodon fourchensis* × *Plethodon ouachitae* 92-110; *Cyprinodon alvarezi* × *Megupsilon aporus* 149-152; *Morone chrysops* × *Morone saxatilis*, *Morone americanus* × *Morone saxatilis* (meristic characters) 513-518.
INCUBATION, *Gehyra mutilata*, *Hemidactylus frenatus*, *Platyurus platyurus* (incubation period) 552-554.
INTRODUCTIONS, *Litoria aurea* (current distribution in New Hebrides) 355-356.
INVASION, *Peprilus burti* (of Atlantic Ocean from Gulf of Mexico) 538-541.
KARYOLOGY, *Anableps dowi* 82-91; *Colisa fasciatus* (somatic G-banded chromosomes and confirmation of female heterogamety) 146-149; *Bufo (americanus, woodhousei woodhousei, w. fawleri, marinus)* (ammoniacal silver staining of nucleolar organizer regions) 341-345; neotropical Cichlidae (41 spp.) (cytotaxonomy) 679-691; genera *Kinosternon*, *Sternotherus*, *Staurotypus* (6 taxa) (banded karyotypes) 692-698.
KEYS (TAXONOMIC), genus *Cophixalus* (4 spp.) 118-121; family Diceratiidae (key to metamorphosed females) 129-144; genus *Microdesmus* (three spp.; eastern Atlantic Ocean) 205; *Kinosternon flavescens* (includes 4 subsp.) 221-222; genus *Rhabdosargus* 702-709.
LARVAE, *Desmognathus fuscus fuscus* (larval feeding behavior) 354; *Rana catesbeiana* (survival rate, population density and development) 447-453; larval fishes (x-radiography) 535-538; *Gobiomorus dormitor* (advantage of schooling behavior) 542-544.
LIFE HISTORY, *Etheostoma blennioides* 191-203.
LIPIDS, *Bufo boreas* (lipid components of prey odors elicit feeding responses) 275-278.
LIVER, *Anguilla rostrata* (phosphate compounds in liver) 360-363.
MAPS (DISTRIBUTION), family Diceratiidae 141.
MERISTICS, *Morone* hybrids (*M. chrysops* × *M. saxatilis*; *M. americana* × *M. saxatilis*) 513-518.
METABOLISM, *Cnemidophorus murinus* (metabolic expenditure and cost of foraging) 573-577; *Xantusia (henshawi, vigilis)*, *Klauberina riversiana*, *Lepidophyma (gaigeae, smithi)* (oxygen consumption) 577-584; *Anolis limifrons* (related to reproduction) 620-626.
MORPHOLOGY, gekkonine and diplodactyline geckos (parallelism and integrated design in foot structure) 1-21; family Diceratiidae (primarily osteology) 129-144; *Gasterosteus aculeatus* (population decline and change in frequency of lateral plates) 635-638; trionychid turtles (packaging problems of head retraction) 655-660.
MOVEMENT, *Peprilus burti* (from Gulf of Mexico into Atlantic Ocean) 538-541.
MUSCLE, *Anguilla rostrata* (phosphate compounds in skeletal muscle) 360-363.
NESTS, *Gasterosteus aculeatus* (nest habitat preference of low-plate number morphs) 525-528.
ODORS, *Bufo boreas* (lipid components of prey odors elicit feeding responses) 275-278.
OLFACTION, *Bufo boreas* (olfactory cues in feeding

- response) 163-165; *Bufo boreas* (feeding responses elicited by lipid components of prey odors) 275-278.
- ONTOGENY, *Ambystoma (maculatum, tigrinum)* (ontogenetic variation in ovum size) 348-350.
- ORGANS, *Plotosus lineatus* (histophysiology of dentritic organs in relation to osmoregulation) 357-360.
- OSMOREGULATION, *Plotosus lineatus* (histophysiology of gills and dentritic organs in relation to osmoregulation) 357-360; *Kinosternon baurii baurii* 548-552.
- OSTEOLOGY, family Diceratiidae 129-144; Ranidae (many spp.) (observations on cranial foramina) 346-348.
- PALEONTOLOGY, fossil gerrhosaur (Miocene of Kenya) 172-174; *Kinosternon arizonense* (Pliocene fossil; determined to be senior synonym of *Kinosternon flavescens stejnegeri*) 175-177; *Libotonius pearsoni* (new species of Percopidae from Eocene of Washington state) 400-405; Nash local herpetofauna (Pleistocene: Aftonian) 747-749.
- PARENTAL CARE, *Desmognathus ochrophaeus* (adaptiveness of care of egg clutches to predation) 332-341.
- PHOTOPERIODICITY, *Cryptobranchus alleganiensis* (activity rhythms) 92-95; *Chrysemys picta*, *Clemmys guttata*, *Sternotherus odoratus* (effect of temperature and photoperiod acclimatization on thermal preferences) 165-169.
- PHYLOGENY, genera *Plethodon* and *Ensatina* (albumin evolution and phylogenetic implications) 502-508.
- PHYSICO-CHEMICAL FACTORS, *Notropis lutrensis* (influence on habitat selection) 70-81.
- PHYSIOLOGY, *Clarias batrachus* (synchronous aerial respiration) 156-158; *Sceloporus cyanogenys* (thermoregulation as related to operant responses for heat at different thermal intensities) 258-266; *Plotosus lineatus* (histophysiology of gills and dentritic organs in relation to osmoregulation) 357-360; *Cnemidophorus murinus* (metabolic expenditure and cost of foraging) 573-577.
- PIGMENTATION, *Poecilia reticulata* (differential response to melanophore-based color mutants) 232-242; *Lacerta dugesii* (dorsal pigmentation found to be related to differential predation pressures rather than to temperature regulation) 250-257.
- POPULATIONS, *Enhydryna schistosa* (population structure) 307-318; *Triturus cristatus* (structure) 350-353; *Rana catesbeiana* (survival rate, population density and development of naturally occurring larvae) 447-453; *Gasterosteus aculeatus* (population decline correlated with change in lateral-plate frequencies) 635-638.
- PREDATION, piscivorous predators and disabled prey 158-160; Brazilian snakes (by mustelid mammals (genus *Galiotis*)) 169-172; *Lacerta dugesii* (dorsal pigmentation found to be related to differential predation pressures than to thermoregulation) 258-266; *Mugil cephalus* (by *Crocodylus niloticus*) 266-269; *Desmognathus ochrophaeus*, *Eurycea bislineata*, *Plethodon (cinereus, jordani, glutinosus)*, *Ambystoma maculatum*, *Gyrinophilus porphyriticus* (effectiveness of anti-predator secretions and behavior against shrew predation) 270-274; *Desmognathus ochrophaeus* (eggs less susceptible to predation when attended by females than when unattended) 332-341; *Salamandra salamandra* (stimulus orientation and stimulus movement pattern in prey-catching behavior) 442-447.
- PRESERVATION, effects of buffered and unbuffered formalin of fish otoliths 155-156.
- PREY, *Damalichthys vocca*, *Embiotoca lateralis* (geographic patterns of prey utilization) 567-572.
- RECORD, *Coryphaenoides thelostomus* (second known specimen) 541-542.
- REPLACEMENT, *Gnathopogon coeruleus* (replacement pattern of pharyngeal teeth) 22-28.
- REPRODUCTION, *Varanus gouldii rosenbergi* 64-69; *Etheostoma blennioides* 201-202; *Enhydryna schistosa* 307-318; *Elaphe (obsoleta, guttata, vulpina)* (reproductive behavior) 319-331; *Melanostigma atlanticum*, *Xenodermichthys copei* (demersal spawning) 363; *Micrurus fulvius tener* (in Texas) 453-463; *Storeria dekayi* (female reproductive biology; Louisiana) 463-466; *Anolis limifrons* (metabolism) 620-626.
- RESPIRATION, *Clarias batrachus* (synchronous aerial respiration) 156-158.
- SALINITY TOLERANCE, *Kinosternon baurii baurii* 548-552.
- SALT GLAND, *Cerberus rhynchops* (possible new salt gland) 661-672.
- SCHOOLING, *Gobiomorus dormitor* (advantage of schooling behavior to fry) 542-544.
- SECRETIONS, *Desmognathus ochrophaeus*, *Eurycea bislineata*, *Plethodon (cinereus, jordani, glutinosus)*, *Ambystoma maculatum*, *Gyrinophilus porphyriticus* (effectiveness of anti-predator secretions and behavior against shrew predation) 270-274.
- SELECTION, *Hyla versicolor* (breeding success related to perch sites rather than size of males) 286-290; numerous amphibian species (sexual selection and sexual dimorphism) 297-306.
- SEX, numerous amphibian species (sexual selection and sexual dimorphism) 297-306.
- SEX DETERMINATION, *Poecilia reticulata* (genetic studies of atypical sex determination) 225-231.
- SEXUAL DIMORPHISM, *Gasterosteus aculeatus* (of integumentary tissues) 533-535.
- SOCIAL, *Python molurus* (social behavior and hierarchy) 466-471.
- SPERM STORAGE, *Acrochordus javanicus* (long term—7 years—sperm storage) 744-745.
- SPAWNING, *Fundulus heteroclitus* (lunar spawning cycle) 291-297; *Melanostigma atlanticum*, *Xenodermichthys copei* (demersal spawning) 363-366; *As-*

- tyanax mexicanus* (breeding method based on annual spawning patterns) 369-371.
- STAINING**, *Bufo* (*americanus*, *woodhousei* *woodhousei*, *w. fowleri*, *marinus*) (ammoniacal silver staining of nucleolar organizer regions) 341-345.
- SURVIVAL**, *Rana catesbeiana* (survival rate of larvae) 447-453.
- SYNONYMY**, gymnotid genus *Hildatia* (junior synonym of *Sternopygus*) 751-752.
- SYSTEMATICS**, *Anableps dowi* (relationships of species to other *Anableps* and of family Anablepidae to other atheriniform fishes) 82-91; *Plethodon* (*ouachitae*, *caddoensis*, *fouchensis*) (genetic relationships as determined by electrophoresis; most closely related to *P. glutinosus*) 95-110; *Cophixalus concinnus* (new species description; relationships; taxonomic key of genus) 118-121; *Cynisca rouxae* (new species; relationship to other congeners) 122-125; *Varicus marilynae* (new species; relationships) 126-128; family Diceratiidae (revision of family; descriptions of two new species) 129-144; *Maxostoma macrolepidotum* (electrophoresis; creatine kinase variability) 152-154; *Kinosternon arizonense* (validity; regarded as subspecies of *K. flavescens*) 175-177; *Etheostoma blennioides* (intraspecific variation, including description of new subspecies) 192-203; *Microdesmus africanus* (new species description; relationships and taxonomic key) 203-205; *Scarus* (*japanensis*, *quoyi*, *iserti*) (senior synonyms or corrected names for *Scarus* (*capistratoides*, *blochii*, *croicensis*, respectively) 206-212; *Kinosternon flavescens* (four subspecies, one new) 212-225; *Rana chiricahuensis* (new species in *Rana pipiens* complex; Arizona) 383-390; *Siganus puelluloides* (new species; Indian Ocean) 390-393; *Rypticus* (*bicolor*, *nigripinnis*, *courtenayi*) (eastern Pacific species) 393-400; *Libotonia pearsoni* (new species of Percopsidae from Eocene of Washington state) 400-405; *Pungitius pungitius* (geographic variation in European populations) 405-412; *Percina* (*roanoka*, *crassa*, *peltata*, *notogramma*) 413-426; *Etheostoma nigrum susanae* 426-430; *Morone* hybrids (*M. chrysops* × *M. saxatilis*; *M. americana* × *M. saxatilis*) (meristic characters) 513-518; *Umbrivaga pyburni* (new species, with comments on genus) 698-701; *Rhambosargus thorpei* (new species; key to genus) 702-709; genus *Urolophus* (review of genus; new species) 714-733; genus *Tubbia* (review of genus) 733-738.
- TAXONOMY**, neotropical Cichlidae (41 spp.) (cytotaxonomy) 679-691.
- TECHNIQUES**, *Bufo* (*americanus*, *woodhousei* *woodhousei*, *w. fowleri*, *marinus*) (ammoniacal silver staining of nucleolar organizer region) 341-345; *Chelydra serpentina* (method to reduce error in weight estimation of freshwater turtles) 346.
- TEETH**, *Gnathopogon coeruleus* (development and replacement pattern of pharyngeal teeth) 22-28.
- TEMPERATURE**, *Alligator mississippiensis* and *Crocodylus acutus* (feeding responses to temperature changes) 48-59; *Gambusia a. affinis* (preferences under field and laboratory conditions) 60-64; *Chrysemys picta*, *Clemmys guttata*, *Sternotherus odoratus* (effect of temperature and photoperiod acclimatization in thermal preferences) 165-169.
- THERMOREGULATION**, *Lacerta dugesii* (dorsal pigmentation shown to be related to differential predation pressures and not to thermal regulation) 250-257; *Sceloporus cyanogenus* (operant responses for heat at different thermal intensities) 258-266.
- TISSUES**, *Gasterosteus aculeatus* (sexual dimorphism of integumentary tissues) 533-535.
- VARIATION**, *Ambystoma* (*maculatum*, *tigrinum*) (ontogenetic variation in ovum size) 348-350.
- VOCALIZATION**, *Anolis grahami* (description and significance) 481-489.
- WATER BALANCE**, *Cerberus rhynchops* (possible new salt gland) 661-672.
- WEIGHT**, *Chelydra serpentina* (method to reduce error in weight estimates of freshwater turtles) 346.
- X-RADIOGRAPHY**, larval and juvenile fishes 535-538.
- ZOOGEOGRAPHY**, ostariophysan zoogeography (alternative hypothesis) 111-118.